



The Kaneka Ubiquinol® (Kaneka QH®) Difference

Made in the USA

Fermented from non-GMO yeast

Subject of 100+ clinical studies

Kosher certified and allergen free

Self-affirmed GRAS

Bioidentical to the body's natural ubiquinol

Compliant with USP monograph

Supporting 200+ brands

Ubiquinol is the active antioxidant form of coenzyme Q10 and the preferred form in biological systems.

Ubiquinol is found in almost every cell of the body.

Ubiquinol supports cellular energy (ATP) production.

Over 95% of the total CoQ10 in the plasma of a healthy individual is in the antioxidant form—ubiquinol.¹



Ubiquinol is a powerful lipid-soluble antioxidant² providing an active defense against oxidative stress, which may affect lipids, proteins, and DNA.³

Ubiquinol is capable of regenerating vitamin E, an essential lipid-soluble antioxidant that can only be obtained from the diet.⁴

Absorption and Bioactivity

2x

Kaneka Ubiquinol® has been shown to be **2x better absorbed than a conventional CoQ10** supplement.⁴

8x

Research demonstrates that 200 mg of Kaneka Ubiquinol® **increases ubiquinol levels by approximately 8x** compared to baseline in healthy adults when taken daily for at least 30 days.⁵

Unlike conventional CoQ10 supplements, **Kaneka Ubiquinol® requires no conversion in the body to perform its antioxidant functions.**^{6,7}

For those already using CoQ10, **switching to Kaneka Ubiquinol® may offer enhanced absorption and bioavailability.**



Formats

Kaneka Ubiquinol® is available as a pure crystalline powder for use in the following product applications:

- Softgels
- Gummies
- Cap-within-cap
- Liposomal formulations
- Liquid capsules
- Other light- and oxygen-controlled environments

Kaneka Ubiquinol® is also available in an air-stable form as Kaneka Q30™, a 30% Ubiquinol powder. Additional applications include:

- Stick packs
- Sachets
- Soft chews

Packaging

- 1 kg or 5 kg units
- MOQ: 1 kg
- Q30 MOQ: 3 kg

Kaneka
NUTRIENTS

Nutrients.Sales@Kaneka.com

KanekaNutrients.com

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Physical & Chemical Properties

Common Name

Ubiquinol

Molecular Formula



Melting Point

Approx. 49.5° C (about 121.1° F)

Stability

4-year shelf life

Appearance


White to pale yellow crystalline powder

Solubility

Water—Practically insoluble

Oil—Soluble

Chemical Structure



Safety

Kaneka Ubiquinol® has a well-established safety profile as demonstrated by extensive clinical trial data.

KANeKA
NUTRIENTS

Nutrients.Sales@Kaneka.com

KanekaNutrients.com

Benefits



Exercise Performance

Kaneka Ubiquinol® supports physical performance during exercise by:

- Promoting a healthy oxidative balance⁸ during exercise
- Supporting energy metabolism⁹
- Supporting cardiovascular performance when training at high altitudes^{10,11}
- Enhancing peak power production in elite athletes when taken at 300 mg/day¹²



Preconception Support

- Ubiquinol supports mitochondrial function and cellular energy generation—essential for oocyte quality and overall egg health.¹³⁻¹⁵
- In an open-label study, Kaneka Ubiquinol® supplementation was associated with improvements in sperm motility of up to 26% and sperm count increases of up to 53% compared to baseline.¹⁶ Other research has reported improvements in morphology up to 24% compared to baseline.¹⁷



Lifelong Wellness

- Supports mitochondrial and cellular function through cellular energy production and antioxidant defense.^{23,29}
- Shown to reduce fatigue at high altitudes.^{10,11}



Menopausal Well-Being

- Kaneka Ubiquinol® supports general health and well-being during and after menopause.^{18,19}
- In a consumer use study, 80% of menopausal women taking 200 mg of Kaneka Ubiquinol® per day reported decreased irritability, sensitivity, stress, and mood swings after 60 days of supplementation.¹⁸



Heart Health

- Improves blood markers associated with heart health.²⁰
- Supports the high energy requirements of the heart.²¹
- Replenishes CoQ10 levels depleted by statin medications.²²
- Benefits vessel health by supporting proper vasodilation, enhancing nitric oxide production, protecting LDL cholesterol from oxidation, and promoting healthy mitochondrial function.⁷



Healthy Aging

- Supplementation with Kaneka Ubiquinol® replenishes plasma ubiquinol levels.⁷
- Research demonstrates a positive correlation between total CoQ10 status and percentage plasma ubiquinol and overall physical functioning in older adults.^{23,24}
- Higher blood ubiquinol levels promote cardiovascular and muscle health.^{7,23}

References

1. Tang PH, et al. HPLC analysis of reduced and oxidized coenzyme Q10 in human plasma. *Clin Chem*. 2001;47(2):256-65.
2. Ernster L, Forsmark-Andrée P. Ubiquinol: an endogenous antioxidant in aerobic organisms. *Clin Invest*. 1993;71(8 Suppl):S60-5.
3. Sies H. Strategies of antioxidant defense. *Eur J Biochem*. 1993;215(2):213-9.
4. Langsjoen PH, Langsjoen AM. Comparison study of plasma coenzyme Q10 levels in healthy subjects supplemented with ubiquinol versus ubiquinone. *Clin Pharmacol Drug Dev*. 2014;3(1):13-7.
5. Hosoe K, et al. Study on safety and bioavailability of ubiquinol (Kaneka QH) after single and 4-week multiple oral administration to healthy volunteers. *Regul Toxicol Pharmacol*. 2007;47(1):19-28.
6. Kubo H, et al. Orally ingested ubiquinol-10 or ubiquinone-10 reaches the intestinal tract and is absorbed by the small intestine of mice mostly in its original form. *J Clin Biochem Nutr*. 2023;72(2):101-6.
7. Sabbatinelli J, et al. Ubiquinol ameliorates endothelial dysfunction in subjects with mild-to-moderate dyslipidemia: a randomized clinical trial. *Nutrients*. 2020;12(4):1098.
8. Sarmiento A, et al. Short-term ubiquinol supplementation reduces oxidative stress associated with strenuous exercise in healthy adults: a randomized trial. *Biofactors*. 2016;42(6):612-22.
9. Martini FH. Metabolism, nutrition and energetics. In: *Fundamentals of Anatomy and Physiology*, 12th ed. Pearson; 2024:943-50.
10. Liu Z, et al. Effect of ubiquinol on electrophysiology during high-altitude acclimatization and de-acclimatization: a substudy of the Shigatse CARdiorespiratory Fitness (SCARF) randomized clinical trial. *Int J Cardiol*. 2024;401:131817.
11. Lv H, et al. Cardiorespiratory fitness and effects of ubiquinol during high-altitude acclimatization and deacclimatization: The SCARF trial. *iScience*. 2025;28(3):112112.
12. Alf D, et al. Ubiquinol supplementation enhances peak power production in trained athletes: a double-blind, placebo controlled study. *J Int Soc Sports Nutr*. 2013;10:24.
13. Zhu Z, et al. Ovarian aging: mechanisms and intervention strategies. *Med Rev* (2021). 2022;2(6):590-610.
14. Zhang M, et al. Coenzyme Q10 ameliorates the quality of postovulatory aged oocytes by suppressing DNA damage and apoptosis. *Free Radic Biol Med*. 2019;143:84-94.
15. Ben-Meir A, et al. Coenzyme Q10 restores oocyte mitochondrial function and fertility during reproductive aging. *Aging Cell*. 2015;14(5):887-95.
16. Thakur AS, et al. Effect of ubiquinol therapy on sperm parameters and serum testosterone levels in oligoasthenozoospermic infertile men. *J Clin Diagn Res*. 2015;9(9):BC01-3.
17. Cakiroglu B, et al. Ubiquinol effect on sperm parameters in subfertile men who have astheno-teratozoospermia with normal sperm concentration. *Nephrourol Mon*. 2014;6(3):e16870.
18. Kaneka Internal Report. Real-life UBIQUINOL study on 200 postmenopausal women. Expansion Consulteam. 2024.
19. Palacios S, et al. Estudio clínico para conocer la eficacia de la coenzima Q-10 sobre la calidad de vida en mujeres postmenopáusicas. *Toko-Gin Pract*. 2019;78(1):3-7. (Proprietary English translation on file.)
20. Onur S, et al. Ubiquinol reduces gamma glutamyl transferase as a marker of oxidative stress in humans. *BMC Res Notes*. 2014;7:427.
21. Becker WM, Deamer DW. Energy from chemical bonds: the aerobic mode. In: *The World of the Cell*. 2nd ed. Benjamin/Cummings; 1991:275-313.
22. Zlatohlavek L, et al. The effect of coenzyme Q10 in statin myopathy. *Neuro Endocrinol Lett*. 2012;33(Suppl 2):98-101.
23. Fischer A, et al. Coenzyme Q10 status as a determinant of muscular strength in two independent cohorts. *PLoS One*. 2016;11(12):e0167124.
24. de la Bella-Garzon R, et al. Levels of plasma coenzyme Q10 are associated with physical capacity and cardiovascular risk in the elderly. *Antioxidants* (Basel). 2022;11(2):279.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.